



REVISTA DE GASTROENTEROLOGÍA DE MÉXICO

www.elsevier.es/rgmx



ORIGINAL ARTICLE

Association of the *Helicobacter pylori* cagA, vacA, and iceA genotypes with chronic follicular gastritis in a Colombian population at high risk for gastric cancer[☆]



Y.H. Carlosama-Rosero^{a,*}, H. Bolaños-Bravo^b, C.H. Sierra-Tórres^c, E.A. Rosero^d

^a Especialista en Anatomía Patológica, Magister en Epidemiología, Grupo Interdisciplinario de Investigación en Salud-Enfermedad, Profesor Universidad Cooperativa de Colombia, Pasto, Colombia

^b Especialista en Anatomía Patológica, Magister en Oncología Molecular, Grupo de Investigación en Genética Humana y Aplicada, Universidad del Cauca, Popayán, Colombia

^c Biólogo, PhD en Epidemiología Molecular, Grupo de Investigación en Genética Humana y Aplicada, Universidad del Cauca, Popayán, Colombia

^d Especialista Salud Ocupacional. Universidad Cooperativa de Colombia, Pasto, Colombia

Received 10 February 2018; accepted 27 March 2018

Available online 6 August 2018

KEYWORDS

Helicobacter pylori;
Gastritis;
Mucosa-associated
lymphoid tissue;
Genotype;
Lymphoma

Abstract

Introduction and objective: Follicular gastritis is associated with *Helicobacter pylori* infection, but little is known of its relation to bacterial genotypes. Our aim was to establish the relation between follicular gastritis and different *H. pylori* strains.

Materials and methods: An analytic case-control study was conducted that included 36 patients with follicular gastritis (cases) and 83 with nonatrophic gastritis (controls). The sociodemographic information was obtained through a questionnaire. Biopsies were evaluated according to the Sydney System and the Wotherspoon scoring system. *Helicobacter pylori* genotyping was performed using the polymerase chain reaction technique. The quantitative variables were presented as mean and standard deviation and the qualitative variables as proportions and absolute frequency. The effect of each variable on outcome (follicular gastritis) was evaluated through the odds ratio and its 95% confidence interval. Statistical significance was set at a $p < 0.05$.

Results: Follicular gastritis was associated with *Helicobacter pylori* infection (OR: 13.41, CI: 1.7-103, $p = 0.01$). The CagA+ genotype was present in 56.5% of the cases and 58% of the controls. The cytotoxic VacAs1m1 strain was present in 82% of the isolates in both groups. IceA1 frequency was 34.8% in the cases and 26% in the controls and the difference was not statistically significant.

[☆] Please cite this article as: Carlosama-Rosero YH, Bolaños-Bravo H, Sierra-Tórres CH, Rosero EA. Asociación de los genotipos cagA, vacA e IceA de *H. pylori* con la gastritis crónica y folicular en una población colombiana con alto riesgo de cáncer gástrico. Revista de Gastroenterología de México. 2019;84:158–164.

* Corresponding author. Tel.: +3217230843; Universidad Cooperativa de Colombia sede Pasto. Calle 18#47-150 Torobajo.
E-mail address: yeharca@hotmail.com (Y.H. Carlosama-Rosero).

PALABRAS CLAVE

Helicobacter pylori;
Gastritis;
Tejido linfóide
asociado a mucosas;
Genotipo;
Linfoma

Conclusions: The population studied had elevated frequencies of cytotoxic *Helicobacter pylori* strains and the iceA1 genotype was more frequent in follicular gastritis.

© 2018 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Asociación de los genotipos cagA, vacA e IceA de *H. pylori* con la gastritis crónica y folicular en una población colombiana con alto riesgo de cáncer gástrico

Resumen

Introducción y objetivo: La gastritis folicular se asocia con la infección por *Helicobacter pylori*; sin embargo, su relación con los genotipos bacterianos es poco conocida. Nuestro objetivo fue establecer la relación entre la gastritis folicular y distintas cepas del *H. pylori*.

Material y métodos: Estudio analítico de casos y controles que incluyó 36 pacientes con gastritis folicular (casos) y 83 con gastritis no atrófica (controles). La información sociodemográfica se obtuvo mediante un cuestionario. Se obtuvieron biopsias que se evaluaron según las clasificaciones de Sydney y Wotherspoon. La genotipificación de *H. pylori* se realizó mediante reacción en cadena de la polimerasa. Para las variables cuantitativas se empleó la media y la desviación estándar. Las variables categóricas se presentan como proporciones y frecuencia absoluta. La OR con su IC del 95% evaluó el efecto de cada variable sobre el desenlace (gastritis folicular). Un valor de $p < 0.05$ se consideró significativo.

Resultados: La gastritis folicular se asoció con la infección por *Helicobacter* OR: 13.41 IC (1.7-103) $p = 0.01$. El genotipo CagA+ estuvo presente en el 56.5% de los casos y 58% de los controles. El 82% de los aislamientos presentó la cepa citotóxica VacAs1m1 en los 2 grupos. La frecuencia de iceA1 fue de 34.8% y 26% en casos y controles respectivamente, diferencias que no fueron estadísticamente significativas.

Conclusiones: Las cepas citotóxicas de *Helicobacter* exhiben elevadas frecuencias en la población estudiada. El genotipo iceA1 fue más frecuente en la gastritis folicular.

© 2018 Asociación Mexicana de Gastroenterología. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction and aims

Follicular gastritis is characterized by a marked mononuclear inflammatory infiltrate and the formation of lymphoid follicles with a germinal center in the lamina propria.^{1,2} For some authors, at least 2 secondary lymphoid follicles should be identified in an area 1 cm from the gastric mucosa to make the diagnosis.³ Disease prevalence is variable. In Japan a prevalence of 0.19% was reported, Greek and French researchers described a prevalence of 12.7% and 14.2%, respectively,^{4,5} and in other studies, prevalence varied from 27.4% to 79.9%.⁶⁻⁸ A Colombian study described prevalence at 8.4%.⁹

The disease is associated with age, *Helicobacter pylori* (*H. pylori*) infection, and the development of mucosa-associated lymphoid tissue (MALT) lymphoma.^{1,10} In relation to age, a greater frequency of the disease has been determined in pediatric patients with persistent dyspeptic symptoms, and in adolescents and young adults. Female sex appears to be a special condition for the development of the pathology.^{5,11} Of greater importance is the relation of follicular gastritis to MALT lymphoma. Even though lymphoma etiology can be linked to multiple factors, *Helicobacter* infection appears to play a dominant role. That role has been demonstrated, following the regression of follicular gastritis and even MALT lymphoma after bacterial eradication.¹² Nevertheless, little is known about the cascade of events

that occurs from follicular gastritis to the appearance of lymphoma.¹³

The relation of bacterial genotypes to certain kinds of diseases is well-known. The cagA+ and vacAs1/m1 genotypes are considered the most virulent strains associated with peptic ulcer, gastric cancer, and severe inflammation,¹⁴ whereas the s2/m2 strain produces limited inflammation. However, little is known about the role of the bacterial genotypes in the origin of follicular gastritis. In that context, the aim of the present study was to establish the relation between the genetic variability of *H. pylori* (genotypes cagA, vacA, and iceA1) and follicular gastritis. The intention was to broaden the understanding of the determining factors of that condition and establish whether there is a bacterial genotype associated with it. Because follicular gastritis is an entity related to gastric lymphoma, the hypothesis is that the most aggressive bacterial genotypes are associated with that type of inflammation.

Materials and methods

An analytic unmatched case-control study was conducted on patients that were admitted to the gastroenterology units of the *Hospital Universitario San José* and *Endovideo* in the city of Popayán (located in Southwest Colombia in the Andes and catalogued as a high-risk area for gastric cancer),¹⁵ within the time frame of January 2008 and December

Table 1 Follicular gastritis severity grade, including the suspicious MALT lymphoma grade.

Grade 0	Normal	Scant inflammatory cells in the lamina propria. No lymphoid follicles
Grade 1	Chronic gastritis	Small clusters of lymphocytes in the lamina propria. No lymphoid follicles
Grade 2	Follicular gastritis	Prominent lymphoid follicles with plasma cells and mantle zone
Grade 3	Suspicious, probably reactive, inflammatory infiltrate	Lymphoid follicles surrounded by small lymphocytes that diffusely infiltrate the lamina propria and occasionally the epithelium
Grade 4	Suspicious infiltrate, probably lymphoma	Lymphoid follicles surrounded by centrocyte-like cells that infiltrate the lamina propria and the epithelium in small groups
Grade 5	MALT lymphoma	Presence of a dense infiltrate of centrocyte-like mononuclear cells in the lamina propria that invade the epithelial lamina—lymphoepithelial lesion—

Source: Modified from Wotherspoon et al.¹⁶

2011. The study sample was obtained through convenience sampling and included patients above 18 years of age, with a histopathologic diagnosis of follicular gastritis or nonatrophic gastritis (the control group). The participants, as well as their parents, had to be born in a municipality of the Cauca department. Patients with atrophy, metaplasia, gastric cancer or other type of cancer, gastric surgery, previous treatment for *H. pylori*, or HIV infection were excluded.

Biologists from the *Grupo de Investigación en Genética Humana y Aplicada (GIGHA)* at the *Universidad del Cauca* applied a questionnaire to collect the sociodemographic variables (age, sex, educational level, income, and ethnic group). Gastric samples were obtained through gastrointestinal endoscopy performed by experienced gastroenterologists. Patients were referred for endoscopy due to dyspeptic symptoms and they arrived for the examination having fasted for at least 8 hours. The participants were not sedated but received a topical solution to anesthetize the pharynx. Five samples were taken: 2 biopsies from the antrum (the greater and lesser curvatures), 2 from the corpus (the greater and lesser curvatures) and one from the incisura angularis. The biopsies were fixed in buffered formalin and stained with the hematoxylin-eosin and Giemsa stains. The histopathologic analysis of each sample was carried out by 2 pathologists. Visual analogue scales were employed according to the Sydney system. The presence of both histologic activity (inflammation due to polymorphonuclear neutrophils) and *H. pylori* was evaluated. Follicular gastritis was defined as the presence of one or more lymphoid follicles with germinal centers in the lamina propria, and its severity was staged on a scale from 1 to 5, according to the Wotherspoon system¹⁶ (Table 1).

To carry out the bacterial genotyping studies, DNA was extracted from the biopsies embedded in paraffin, utilizing the Chelex technique (No. C7901; Sigma, St. Louis, MO). The *cagA*, *iceA1*, and *vacA* genes were amplified using the polymerase chain reaction (PCR) method previously described by Yamaoka et al.¹² The PCR mixtures were prepared with 50 ng of genomic DNA, 100 μ mol dNTPs, 2.5 μ l of 10X PCR buffer, 1.0 mM MgCl₂, 1 U of Taq DNA polymerase (No. M1665; Promega, Madison, WI), and 30 pmol of each of the primers, as indicated in Table 2. The reactions began with a denaturalization step at 95 °C for 1 minute, followed by 35 denaturalization cycles at 94 °C for 1 minute, hybridization at 52 °C for 1 minute and extension at 72 °C for 1 minute, and a final extension step at 72 °C for 10 minutes. The products were analyzed through electrophoresis in 2% agarose gels at

80 volts for 40 minutes and the genotypes were recognized according to the expected size of the base pairs.

The NCTC-11637 and NCTC-11638 *H. pylori* strains and the clinical 3062 isolate were provided by the *Instituto Nacional de Cancerología de Colombia* to be used as positive controls. In addition, all the PCR tests had internal amplification controls and adequate molecular markers (ladders).

Demographic information biases were controlled through a questionnaire with closed questions applied by members of the GIGHA, who received instructions and guidance from gastroenterologists and pathologists. Histopathologic evaluation was validated by a second pathologist.

The molecular biology tests were performed according to worldwide accepted protocols. Pilot tests were carried out to verify the quality of the reagents and the extraction kits and the laboratory equipment was previously calibrated. Cases of bacterial coinfection determined through the presence of more than one genotype were excluded from the analysis to prevent interpretation bias of the genetic information and to calculate the true effect of each genotype on the response variable.

Because the variable of age had normal distribution through the Kolmogorov-Smirnov test, it was presented as mean and standard deviation. The differences between the means were evaluated using the Student's t test, whereas the differences in proportions were evaluated through the chi-square test of independence. The OR and its p value were used to evaluate the effect of each study variable on the response variable (follicular gastritis). A multivariate logistic regression analysis was carried out, with sex and age as possible confounders. Statistical significance was set at a $p < 0.05$. The statistical analysis was performed using the SPSS version 23 program.

The participants agreed to partake in the study and signed statements of informed consent. The study was approved by the Ethics Committee for Scientific Research of the *Universidad del Cauca*.

Results

From a total of 808 participants selected within the study time frame, 119 patients met the inclusion criteria. Of those patients, 83 corresponded to the 70% that had nonatrophic gastritis and 36 to the 30% diagnosed with follicular gastritis. The age range in the patients with nonatrophic gastritis was 19 to 81 years, with a mean of 44.11 years. The age range in the patients with follicular gastritis was 21 to 82 years, with

Table 2 Sequence of primers for amplification through PCR.

Genes and regions	Sequence (5'→3')	Size (bp)
cagA	GATAACAGGCAAGCTTTTGAGG CGTCAAAAAGATTGTTTGGCAGA	349
iceA1	GTGTTTTTAACCAAAGTATC CTATAGCCASTYTCTTTGCA	247
vacA s1/s2	ATGGAAATACAACAAACACAC CTGCTTGAATGCGCCAAAC	259/286
vacA m1/m2	CAATCTGTCCAATCAAGCGAG GCGTCTAAATAATTCCAAGG	570/645

Table 3 Sociodemographic characteristics of the patients.

Variables		N = 119		OR	CI	p value
		Chronic gastritis n (%)	Follicular gastritis n (%)			
Coded age	Greater than or equal to 40 years	47 (56.6)	17 (47.2)	1	1	1
Sex	Under 40 years	36 (43.4)	19 (52.8)	1.45	(0.66-3.2)	0.34
	Male	56 (67.5)	24 (66.7)	1	1	1
Educational level	Female	27 (32.5)	12 (33.3)	1.03	(0.45-2.38)	0.93
	Technical or higher education	28 (33.7)	8 (22.2)	1	1	1
Income category	None to middle school	55 (66.3)	28 (77.8)	1.78	(0.71-4.41)	0.21
	> a 1 MMW	40 (48.2)	17 (47.2)	1	1	1
Ethnic group	< a 1 MMW	43 (51.8)	19 (52.8)	1.03	(0.47-2.2)	0.92
	Amerindian	1 (1.2)	2 (5.6)	1	1	1
	White	5 (6)	0 (0)	NA	NA	NA
	Black	1 (1.2)	6 (16.7)	3	(0.12-73)	0.5
	Mestizo	76 (91.6)	27 (75)	0.17	(0.01-2.4)	0.16
	Mulatto	0 (0)	1 (2.8)	NA	NA	NA

MMW: monthly minimum wage; NA: not applicable.

P values < 0.05 are statistically significant.

a mean of 41.86 years. The Student's t test showed no significant differences in the comparison of age means ($p=0.48$). In relation to ethnicity, the large majority of patients were mestizos, but there were 6 (17%) black patients in the follicular gastritis group. No statistically significant differences in the sociodemographic variables were found (Table 3).

The relation between *H. pylori* infection and the follicular inflammatory response was evaluated in the histopathologic study. Thirty-five patients (97.2%) in the follicular gastritis group had the infection, whereas only 60 (72.3%) patients in the control group presented with it. There was a statistically significant association between infection and follicular gastritis; OR: 13.41, IC (1.7-103) $p=0.01$.

A statistically significant association also was found in the evaluated relation between bacterial infection and inflammatory activity (OR: 337.33, IC: [53.1-2142], $p=0.00$).

Follicular gastritis was classified as Wotherspoon grade 2 gastritis in the majority of the case group patients (23/36), as grade 3 gastritis in 12 of the case patients (33%), and as grade 4 gastritis in only one case patient. By sex, 6 men and 17 women had grade 2 gastritis, 6 men and 6 women

presented with grade 3, and one woman, 42 years old, had grade 4. There were no documented cases of MALT lymphoma.

Of the total of 119 participants, 46 patients were excluded from the genetic analysis: more than one vacA bacterial genotype (coinfection) was found in 20 patients and gene amplification was not achieved in 26 patients (Table 4).

Through molecular testing, the *H. pylori* of 73 participants (50 patients with nonatrophic gastritis and 23 patients with follicular gastritis) was characterized. Regarding the distribution of the strains, 29 (58%) patients in the control group were positive for the cagA genotype, whereas in the follicular gastritis group that genotype was described in 13 (56.5%) patients. No statistically significant differences were found in relation to the cagA genotype (the OR for developing follicular gastritis in cagA+ patients was 0.79, CI=0.29-2.13, and $p=0.64$).

In the patients with follicular gastritis, 8 (34.8%) were iceA1 positive, whereas in the control group that genotype was found in 13 (26%) patients. Those differences were not statistically significant (OR: 1.83, CI: 0.64-5.22, $p=0.26$). The s1, s2, m1, and m2 subtypes were determined in the

Table 4 Genetic information not included in the analysis due to bacterial coinfection and absence of amplification.

	Nonatrophic gastritis N = 83 (%) [*]	Follicular gastritis N = 36 (%) [*]
Coinfection	14 (16.9)	6 (16.7)
Not amplified by PCR	19 (22.9)	7 (19.4)

PCR: polymerase chain reaction.

^{*} Percentage within each study group.

Table 5 Distribution of the *cagA*, *iceA1*, and *vacA* genotypes in the study groups; n = 73.

Genotype	Chronic gastritis, n (%)	Follicular gastritis, n (%)	OR	CI	p value
CagA-/IceA-	17 (34)	8 (34.8)	1	1	1
CagA-/IceA+	4 (8)	2 (8.7)	1.06	0.16-7.06	0.95
CagA+/IceA-	20 (40)	7 (30.4)	0.74	0.22-2.47	0.63
CagA+/IceA+	9 (18)	6 (26.1)	1.41	0.37-5.38	0.61
VacA-s2m2	7 (14)	1 (4.3)	1	1	1
VacA-s1m2	2 (4)	3 (13)	10.5	0.67-165.1	0.09
VacA-s1m1	41 (82)	19 (82.6)	3.24	0.37-28.26	0.07

P values < 0.05 are statistically significant.

vacA bacterial genotype analysis and they were grouped into 4 categories, according to their virulent profile. Those groups were then related to the other genotypes (*iceA1* and *cagA*). The *cagA*-/*iceA1*- genotypes and the *vacA* s2/m2 subtypes were selected as contrast groups. The analysis showed a high prevalence (82%) of the *vacA* s1/m1 subtypes for the 2 patient groups with no significant differences (OR: 1.04, CI: 0.28-3.81, p = 0.95).

The s2/m1 subtypes were not documented in either of the 2 groups and the analysis of the rest of the subtypes produced no statistically significant differences (Table 5).

Even though the bivariate analysis showed no statistically significant associations between follicular gastritis and the age and sex variables, a review of the literature revealed a probable relation to the response variable. Therefore, the possible confounding effect of those variables was examined in a logistic regression model that included *H. pylori* infection (Table 6).

Discussion and conclusion

The relation between follicular gastritis and age has been referenced by various authors.^{3,8,17} Even though a young adult population would be expected in our study (given that patients with the advanced age-related conditions of atrophy, metaplasia, dysplasia, and cancer were excluded),¹⁸ the age range of the participants was from 19 to 82 years. Our results showed that, in general terms, the population with follicular gastritis was younger than the patients with chronic gastritis. That finding is similar to results published by other authors.^{19,20}

The association between patterns of inflammatory response and ethnicity has been reported as an important risk factor for or protective factor against different diseases, including gastric cancer.²¹⁻²³ For example, the risk for gastric cancer and gastric inflammation grade have recently

been proposed to be a product of a phenomenon of immunologic tolerance that is dependent on the population ancestry and the bacterial ancestry of *H. pylori*.²² Our study participants, as well as their parents, were born in the Cauca department, and the majority of the patients with follicular gastritis were mestizo. Given the small number of patients in the present study, our conclusions are limited, but that finding is interesting, since it suggests the possibility that population ancestry and ethnic origin could influence the follicular inflammation pattern, something that has not been described in the literature.

The prevalence of *Helicobacter* infection in patients with follicular gastritis varies between 68% and 100%, in accordance with the number of biopsies taken.^{9,24} The Sydney protocol was applied in the present study, which stipulates taking 5 biopsies, enabling significant sampling of the gastric mucosa and a reliable histologic determination.²⁵ Infection was detected through microscopic study in 97.2% of the patients with follicular gastritis, showing a significant difference with the control group (OR: 13.41, CI: 1.7-103, p = 0.01). That relation even persisted in the multivariate model that included the confounding variables. Likewise, a significant association was shown between inflammatory activity and the presence of bacterial infection. The link between infection and active follicular gastritis is widely recognized in the medical community and has been documented by different researchers.^{3,5,26,27}

The role of bacterial genotypes in relation to MALT lymphoma has been evaluated in different studies. In their analyses, Eck, Ye, and Koehler, et al., were able to establish statistically significant associations between lymphoma and the *cagA*, *vacA*s1/m1, and *iceA1* genotypes.²⁸⁻³⁰ However, there are few studies that relate bacterial genotypes and follicular gastritis. Specifically, Soltermann et al. evaluated the presence of lymphoid follicles in the antrum and corpus and correlated the information with the *cagA* and *vacA* alleles and genotypes, demonstrating no sig-

Table 6 Multivariate model adjusted by categorized age and by sex. The measures of association of the variables of interest in relation to outcome (follicular gastritis) are shown.

Variable	Crude OR	CI	p value	Adjusted OR	CI	p value
Age under 40 years	1.45	0.66-3.2	0.34	1.26	0.44-3.62	0.66
Female sex	1.03	0.45-2.38	0.93	1.27	0.43-3.8	0.66
<i>Helicobacter pylori</i>	13.41	1.7-103	0.01	8.7	1.1-70.9	0.04

P values < 0.05 are statistically significant.

nificant relation.³¹ Those findings were similar to ours. Other descriptive and analytic studies have been conducted in Latin America that show associations between bacterial genotypes and the pathologies of ulcer, cancer, and gastritis.³²⁻³⁵ To the best of our knowledge, the present study is the first to examine the relation of bacterial genotypes to follicular gastritis in a Colombian population.

In Colombia, Quiroga and Citelly reported cagA genotype frequencies of 50% and 51.4% in patients with gastritis,^{35,36} figures similar to that found in the present study (58% in the controls).

In relation to the vacAs1/m1 genotypes in the study by Citelly et al., they showed a prevalence of 43% for the s1 strain and of 40% for the m1 strain, whereas prevalence in our findings was above 80%. That information signifies that our study population had more aggressive bacterial genotypes, even at early gastric disease stages. It should be emphasized that none of the patients in the present study had atrophy, metaplasia, or dysplasia, and that the majority of patients in the case group presented with grade 2 or 3 follicular gastritis. In accordance with the knowledge there is on the cascades involved in carcinogenesis proposed by Pelayo Correa, and with the Wotherspoon model,^{16,37} the histologic findings in our study population would suggest a low risk for the development of cancer (both intestinal adenocarcinoma and MALT lymphoma), but the presence of cytotoxic strains would make a more careful endoscopic and histopathologic follow-up of our patients a necessity. Those genotypic differences would explain, at least partially, why patients from the Cauca department have higher incidence rates of cancer, compared with other areas of the country.¹⁵

The iceA1 genotype is considered by some authors to be a virulence factor involved in the cellular adhesion and development of ulcers, whereas others believe its prevalence depends on the geographic distribution of the bacterium.³⁸⁻⁴⁰ In the present study, there was no significant change in the distribution of the iceA1 genotype between the groups compared, but it was more frequent in the follicular gastritis group than in the nonatrophic gastritis group. Those data suggest a probable relation between the follicular inflammatory response and the iceA1 genotype.

The reduced number of data included for the genetic analyses was one of the limitations of our study. The verification of pH in the formalin utilized for biopsy collection or the use of transport means and culture methods as alternatives for facilitating bacterial DNA recovery are suggested for future studies. We hope to motivate the scientific community to evaluate the role of the iceA1 genotype, using larger sample sizes.

In conclusion, our study results showed that follicular gastritis and inflammatory activity are reliable markers for the

presence of *H. pylori* infection and that the genotypes with a high oncogenic risk predominated in the study population. Therefore, its diagnosis should alert the clinician, so that progression to diseases with greater morbidity and mortality is prevented. In addition, it was suggested that the iceA genotype could play a role in the development of follicular gastritis.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients and/or subjects mentioned in the article.

Financial disclosure

The present study was financed by the *Programa de Salud de Colciencias*, project code 1103-519-29123.

Conflict of interest

The authors of the present article declare that there is no conflict of interest.

Acknowledgements

The authors wish to thank the patients for their participation in the study and the personnel of the *Hospital Universitario San José* and *Endovideo* for their invaluable support. In addition, they wish to thank the *Laboratorio de Genética Humana* and the *Departamento de Patología* of the *Facultad de Ciencias de la Salud* of the *Universidad del Cauca* for their support in carrying out this project.

References

1. Genta RM, Hamner HW, Graham DY. Gastric lymphoid follicles in *Helicobacter pylori* infection: Frequency, distribution, and response to triple therapy. *Human Pathol.* 1993;24:577-83.
2. De Giacomo C, Fiocca R, Villani L, et al. *Helicobacter pylori* infection and chronic gastritis: Clinical, serological, and histologic correlations in children treated with amoxicillin and

- colloidal bismuth subcitrate. *J Pediatr Gastroenterol Nutr.* 1990;11:310–6.
3. Ma Z, Tanizawa T, Nihei Z, et al. Follicular gastritis associated with *Helicobacter pylori*. *J Med Dent Sci.* 2000;47:39–47.
 4. Ladas SD, Rokkas T, Georgopoulos S, et al. Predictive factors and prevalence of follicular gastritis in adults with peptic ulcer and nonulcer dyspepsia. *Dig Dis Sci.* 1999;44:1156–60.
 5. Zerbib F, Vialette G, Cayla R, et al. Follicular gastritis in adults. Relations with *Helicobacter pylori*, histological and endoscopic aspects. *Gastroenterologie clinique et biologique.* 1993;17:529–34.
 6. Eidt S, Stolte M. Prevalence of lymphoid follicles and aggregates in *Helicobacter pylori* gastritis in antral and body mucosa. *J Clin Pathol.* 1993;46:832–5.
 7. Genta RM, Hamner HW. The significance of lymphoid follicles in the interpretation of gastric biopsy specimens. *Arch Pathol Lab Med.* 1994;118:740–3.
 8. Chen X, Liu W, Shi Y, et al. *Helicobacter pylori* associated gastric diseases and lymphoid tissue hyperplasia in gastric antral mucosa. *J Clin Pathol.* 2002;55:133–7.
 9. Marín JDM, Riveros SCH. Hiperplasia linfocitaria folicular gástrica e infección por *Helicobacter pylori* en adultos colombianos. *Rev Colomb Gastroenterol.* 2009;24:148–56.
 10. Zullo A, Hassan C, Ridola L, et al. Gastric MALT lymphoma: Old and new insights. *Ann Gastroenterol.* 2014;27:27–33.
 11. Bahú MGS, da Silveira TR, Maguilnick I, et al. Endoscopic nodular gastritis: An endoscopic indicator of high-grade bacterial colonization and severe gastritis in children with *Helicobacter pylori*. *J Pediatr Gastroenterol Nutr.* 2003;36:217–22.
 12. Morgner A, Lehn N, Andersen LP, et al. *Helicobacter heilmannii*-associated primary gastric low-grade MALT lymphoma: Complete remission after curing the infection. *Gastroenterology.* 2000;118:821–8.
 13. Thiede C, Morgner A, Alpen B, et al. What role does *Helicobacter pylori* eradication play in gastric MALT and gastric MALT lymphoma? *Gastroenterology.* 1997;113:61–4.
 14. Blaser MJ. Science, medicine, and the future: *Helicobacter pylori* and gastric diseases. *Br Med J.* 1998;316:1507–10.
 15. Pardo Ramos C, Cendales Duarte R. Incidencia estimada y mortalidad por cáncer en Colombia 2002–2006. Incidencia estimada y mortalidad por cáncer en Colombia 2002–2006. Instituto Nacional de Cancerología; 2010. p. lil-1668548.
 16. Wotherspoon A, Doglioni C, Isaacson P. Low-grade gastric B-cell lymphoma of mucosa-associated lymphoid tissue (MALT): A multifocal disease. *Histopathology.* 1992;20:29–34.
 17. Raymond J, Bergeret M, Benhamou P, et al. A 2-year study of *Helicobacter pylori* in children. *J Clin Microbiol.* 1994;32:461–3.
 18. Martínez JD, Henao SC, Granados C. The corporal atrophic chronic gastritis and the age. *Rev Colomb Gastroenterol.* 2007;22:17–22.
 19. Zaitoun A. The prevalence of lymphoid follicles in *Helicobacter pylori* associated gastritis in patients with ulcers and non-ulcer dyspepsia. *J Clin Pathol.* 1995;48:325–9.
 20. Önal IK, Sökmensüer C, Önal ED, et al. Clinical and pathological features of nodular gastritis in adults. *Turkish J Med Sci.* 2009;39:719–23.
 21. Lee I, Lee H, Kim M, et al. Ethnic difference of *Helicobacter pylori* gastritis: Korean and Japanese gastritis is characterized by male-and antrum-predominant acute foveolitis in comparison with American gastritis. *World J Gastroenterol.* 2005;11:94–8.
 22. Kodaman N, Pazos A, Schneider BG, et al. Human and *Helicobacter pylori* coevolution shapes the risk of gastric disease. *Proceed Nat Acad Sci.* 2014;111:1455–60.
 23. Segal I, Ally R, Sitas F, et al. *Helicobacter pylori*: The African enigma. *Gut.* 1998;43:300–1.
 24. Yakoob MY, Hussain AS. Chronic gastritis and *Helicobacter pylori*: A histopathological study of gastric mucosal biopsies. *J Coll Phys Surg Pakistan.* 2010;20:773–5.
 25. Bravo LE, Bravo JC, Realpe JL, et al. Fuentes de variabilidad en el diagnóstico de gastritis atrófica multifocal asociada con la infección por *Helicobacter pylori*. *Colombia Médica.* 2008;39:58–65.
 26. Mejía C, Vera C, Huiza-Espinoza L. Association between follicular gastritis and *Helicobacter pylori* in children seen at a public hospital in Peru. *Rev Gastroenterol México (English ed).* 2016;81:80–5.
 27. Wotherspoon A, Ortiz Hidalgo C, Falzon M, et al. *Helicobacter pylori*-associated gastritis and primary B-cell gastric lymphoma. *The Lancet.* 1991;338:1175–6.
 28. Koehler C, Mues M, Dienes H, et al. *Helicobacter pylori* genotyping in gastric adenocarcinoma and MALT lymphoma by multiplex PCR analyses of paraffin wax embedded tissues. *Molecular Pathol.* 2003;56:36–42.
 29. Eck M, Schmausser B, Haas R, et al. MALT-type lymphoma of the stomach is associated with *Helicobacter pylori* strains expressing the CagA protein. *Gastroenterology.* 1997;112:1482–6.
 30. Ye H, Liu H, Attygalle A, et al. Variable frequencies of t (11;18)(q21; q21) in MALT lymphomas of different sites: Significant association with CagA strains of *H pylori* in gastric MALT lymphoma. *Blood.* 2003;102:1012–8.
 31. Soltermann A, Koetzer S, Eigenmann F, et al. Correlation of *Helicobacter pylori* virulence genotypes vacA and cagA with histological parameters of gastritis and patient's age. *Modern Pathol.* 2007;20:878–83.
 32. Yamaoka Y, Kodama T, Gutierrez O, et al. Relationship between *Helicobacter pylori* iceA, cagA, and vacA status and clinical outcome: Studies in four different countries. *J Clin Microbiol.* 1999;37:2274–9.
 33. Brito CA, Silva L, Jucá N, et al. Prevalence of cagA and vacA genes in isolates from patients with *Helicobacter pylori*-associated gastroduodenal diseases in Recife, Pernambuco, Brazil. *Memorias do Instituto Oswaldo Cruz.* 2003;98:817–21.
 34. Paniagua GL, Monroy E, Rodriguez R, et al. Frequency of vacA, cagA and babA2 virulence markers in *Helicobacter pylori* strains isolated from Mexican patients with chronic gastritis. *Annals Clin Microbiol Antimicrob.* 2009;8:14.
 35. Trujillo E, Martínez T, Bravo MM. Genotyping of *Helicobacter pylori* virulence factors vacA and cagA in individuals from two regions in Colombia with opposing risk for gastric cancer. *Biomédica.* 2014;34:567–73.
 36. Cittelly D, Huertas M, Martínez J, et al. *Helicobacter pylori* genotypes in non atrophic gastritis are different of the found in peptic ulcer, premalignant lesions and gastric cancer in Colombia. *Rev Med Chile.* 2002;130:143–51.
 37. Correa P. Human gastric carcinogenesis: A multistep and multifactorial process—first American Cancer Society award lecture on cancer epidemiology and prevention. *Cancer Res.* 1992;52:6735–40.
 38. Quiroga AJ, Diana Marcela C, Bravo MM. BabA2, oipA and cagE *Helicobacter pylori* genotypes in Colombian patients with gastroduodenal diseases. *Biomédica.* 2005;25:325–34.
 39. Caner V, Yilmaz M, Yonetcı N, et al. *H pylori* iceA alleles are disease-specific virulence factors. *World J Gastroenterol.* 2007;13:2581–5.
 40. Höcker M, Hohenberger P. *Helicobacter pylori* virulence factors—one part of a big picture. *The Lancet.* 2003;362:1231–3.