



CLINICAL IMAGE IN GASTROENTEROLOGY

## Todani type IVa choledochal cyst<sup>☆</sup>

### Quiste de colédoco tipo IVa de Todani

R. Soto-Solis\*, L.A. Waller



*Servicio de Endoscopia, Centro Médico Nacional 20 de Noviembre, Instituto de Seguridad y Servicios Sociales de los Trabajadores del Estado (ISSSTE), Mexico City, Mexico*


A 39-year-old woman was referred to our hospital center for the diagnostic evaluation of asymptomatic cholestasis. Nuclear magnetic resonance cholangiography (fig. 1) and endoscopic retrograde cholangiopancreatography (ERCP) (figs. 2 and 3) were carried out. Both studies revealed marked cystic dilation of the extrahepatic and intrahepatic bile ducts and a normal intrapancreatic bile duct. The ERCP identified a common channel of the pancreatic duct and the common bile duct (fig. 2). The incidence of congenital choledochal cysts is 1/100,000 to 1/150,000, and the most widely accepted cause is an anomalous junction of the pancreatic duct and the common bile duct.




**Figure 1** NMR cholangiography showing cystic dilation of the intrahepatic bile ducts and the proximal extrahepatic bile ducts.

<sup>☆</sup> Please cite this article as: Soto-Solis R, Waller LA. Quiste de colédoco tipo IVa de Todani. *Revista de Gastroenterología de México*. 2019;84:400–401.

\* Corresponding author. Camino a Santa Teresa 1055 Torre Ángeles Consultorio 650, Colonia Héroes de Padierna, Delegación Magdalena Contreras, CP 10700, Mexico City, Mexico. Tel.: +52(55)1368-9214  
E-mail address: [rodrigosotomd@hotmail.com](mailto:rodrigosotomd@hotmail.com) (R. Soto-Solis).



**Figure 2** ERCP showing the cystic dilation of the intrahepatic bile ducts and the proximal extrahepatic bile ducts. Note the common channel of the pancreatic duct and the common bile duct.



**Figure 3** ERCP showing the intrahepatic cystic dilation. Note the opening of the posterior right hepatic duct into the common bile duct.