



REVISTA DE GASTROENTEROLOGÍA DE MÉXICO

www.elsevier.es/rgmx



ORIGINAL ARTICLE

Prevalence of and factors associated with *Helicobacter pylori* infection in preschoolers in Havana, Cuba: A population-based study[☆]



S.J. Venero-Fernández^{a,*}, I. Avila-Ochoa^b, L. Menocal-Herredia^a,
Y. Caraballo-Sánchez^a, F.M. Rosado-García^a, R. Suárez-Medina^a, P. Varona-Pérez^a,
A.W. Fogarty^c, National History of Wheezing in Cuba (HINASIC) Study Group[◇]

^a Instituto Nacional de Higiene, Epidemiología y Microbiología, La Habana, Cuba

^b Departamento de Gastroenterología, Hospital Universitario Pediátrico Docente Centro Habana, La Habana, Cuba

^c Nottingham Biomedical Research Unit, Division of Epidemiology and Public Health, University of Nottingham, UK

Received 19 April 2018; accepted 25 March 2019

Available online 14 April 2020

KEYWORDS

Helicobacter pylori;
Prevalence;
Risk factors;
Cuba

Abstract

Introduction: *Helicobacter pylori* (*H. pylori*) infection is usually acquired in childhood. It has not been widely studied in individuals within the pediatric age range in Cuba.

Aim: To identify the prevalence of the infection and its associated risk factors in 3-year-old children in Havana.

Materials and methods: An analytic, cross-sectional, epidemiologic analysis was conducted on 1,274 children belonging to the cohort of participants in the Natural History of Wheezing in Cuba study (HINASIC for its Spanish acronym) that were 3 years of age and provided a stool sample. *H. pylori* infection was identified by determining the *H. pylori* antigen (Ag) in stool, utilizing the commercial Spinreact kit, from Spain. The data were collected through a questionnaire applied by the researchers that included sociodemographic, environmental, and lifestyle variables, as well as infection from other parasites. Prevalence and the prevalence ratio with a 95% confidence interval were calculated and the dichotomous logistic regression analysis was employed.

Results: The prevalence of positive *H. pylori* Ag was 5%. Sleeping together was the risk factor found (PR: 1.27; 95% CI: 1.03-1.50). Protective factors were drinking water from water delivery trucks (PR: 0.16; 95% CI: 0.04-0.72) and living in a nuclear family unit (PR: 0.94; 95% CI: 0.85-0.99).

[☆] Please cite this article as: Venero-Fernández SJ, Avila-Ochoa I, Menocal-Herredia L, Caraballo-Sánchez Y, Rosado-García FM, Suárez-Medina R, et al. Prevalencia y factores asociados a infección por *Helicobacter pylori* en preescolares de La Habana, Cuba. Estudio de base poblacional. Revista de Gastroenterología de México. 2020;85:151–159.

* Corresponding author at: Dirección: Infanta No 1158 e/Llinás y Clavel, Código Postal 10300, La Habana, Cuba. Tel.: (537) 870 5531-34 ext 141.

E-mail address: silviavf@infomed.sld.cu (S.J. Venero-Fernández).

[◇] List of collaborating researchers at the end of the article.

PALABRAS CLAVE

Helicobacter pylori;
Prevalencia;
Factores de riesgo;
Cuba

Conclusions: The prevalence of *H. pylori* infection in early childhood places Havana in an intermediate position at the international level. To control the infection, causal studies should be conducted and opportune interventions implemented.

© 2020 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Prevalencia y factores asociados a infección por *Helicobacter pylori* en preescolares de La Habana, Cuba. Estudio de base poblacional

Resumen

Introducción y objetivo: La infección por *Helicobacter pylori* (*H. pylori*) es usualmente adquirida en la infancia. En Cuba su estudio en las edades pediátricas es un campo poco explorado.

Objetivo: Identificar la prevalencia de la infección y los factores asociados en niños de 3 años de edad de La Habana.

Material y métodos: Se realizó un estudio epidemiológico transversal analítico con 1274 niños de 3 años de edad, que aportaron muestra de heces, provenientes de la cohorte de nacimientos de La Habana (HINASIC). Se identificó la infección por *H. pylori*, utilizando el paquete para determinación de antígeno (Ag) de *H. pylori* en heces de la casa comercial Spinreact, España. La recolección de datos fue a través de un cuestionario administrado por los investigadores que incluyen variables sociodemográficas, ambientales, estilo de vida e infección por otros parásitos. Se calcularon prevalencias, razón de prevalencia con intervalos de confianza de 95% y regresión logística dicotómica.

Resultados: La prevalencia de Ag de *H. Pylori* positivo fue de 5%. Dormir acompañado fue el factor de riesgo encontrado RP 1.27 (IC 95%: 1.03-1.50). Consumo de agua de pipa RP 0.16 (IC 95%: 0.04-0.72) y vivir en un núcleo familiar con ambos padres RP 0.94 (IC 95%: 0.85-0.99) fueron factores protectores.

Conclusiones: La prevalencia de infección de *H. pylori* en la infancia temprana ubica a La Habana internacionalmente en una posición intermedia. Estudios de causalidad e intervenciones futuras deberán ser tenidos en cuenta para el control de la infección.

© 2020 Asociación Mexicana de Gastroenterología. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction and aims

Helicobacter pylori (*H. pylori*) colonization of the stomach is the most common of the chronic bacterial infections in humans.¹ It is considered one of the main findings in gastroenterology and is one of the microorganisms of greatest interest in human pathology.^{2,3} Its origin dates back some 58,000 years.⁴ Between 1985-89, Warren and Marshall⁵ associated the presence of the bacterium with chronic gastritis and ulcer, and since then, much work has been carried out in that field. The genesis of chronic gastritis, duodenal and gastric ulcer in children and adults, and both gastric adenocarcinoma and gastric mucosa-associated lymphoid tissue (MALT) lymphoma have been attributed to it. Once acquired, the infection persists, and the majority of infected individuals remain asymptomatic.

Calculations suggest that approximately half of the world's population is colonized by *H. pylori*, with wide variability between different populations, possibly related to different factors, such as the virulence of the microorganism, host susceptibility, antibiotic use, and environmental conditions that include socioeconomic level. Genetic,

racial, and cultural factors are less clearly involved.^{6,7} Transmission via the digestive route is well-accepted.⁶ Prevalence ranges from 1-2% in studies conducted on asymptomatic children in the Netherlands, to over 90% in Bangladesh.⁶ In developing countries, children are infected mainly before 10 years of age, and the highest prevalence peak in adults (80%) is before 50 years of age. In developed countries, studies show that infection is rarely produced before 10 years of age. It increases to 10% between 18 and 30 years of age and to 50% above 60 years of age.⁸

The prevalence of *H. pylori* infection in Cuba has not been widely analyzed. The majority of studies have been conducted on pediatric and adult populations with gastrointestinal symptoms. A high rate of *H. pylori* infection has been identified in adult patients with gastric ulcer (91%), duodenal ulcer (99%), and chronic gastritis (94%), with an elevated expression of the *cagA*⁺ gene.⁹⁻¹² There are fewer studies on children, but a relation to anemia in school children (79%)¹³ and to diagnoses of antral chronic gastritis (67.7%) has been observed.¹⁴ Opportune preventive measures for stopping *H. pylori* transmission will reduce the incidence of those diseases and will depend on knowledge

of the history of *H. pylori* infection. Therefore, the aim of the present study was to identify the prevalence of *H. pylori* infection and the associated socioeconomic, environmental, and lifestyle factors in the pediatric population of 3-year-old children in the province of Havana.

Materials and methods

Study population

The participants of the present analysis belong to the third year of the Natural History of Wheezing in Cuba (*HINASIC* for its Spanish acronym) study, a prospective, longitudinal, population study on children, whose aim was to identify risk factors for the development of asthma. Detailed information on the methodological design of the study, including sample selection, and the inclusion and exclusion criteria were previously described.¹⁵ A total of 1,543 three-year-old children who attended any of the 17 polyclinics of the 4 municipalities of Havana (*Arroyo Naranjo, Cerro, Habana del Este, La Lisa*) were interviewed within the time frame of March 2012 to March 2013. Of those children, 1,274 (83%) were eligible to participate in the present study by providing a stool sample.

Data collection

The data were obtained through the application of a questionnaire administered by an interviewer (pediatricians or family physicians) to parents or guardians, after they had given their informed consent for their children to participate in the study. Information on demographics, lifestyle, and home environment, as well as specific questions on anthropometric measurements (weight and height) at the time of the interview, were collected. *H. pylori* infection was diagnosed from the stool sample provided by the participants, utilizing the commercial Spinreact kit (<http://www.spinreact.com>). As stated in the technical label, it is a qualitative immunochromatographic test that has 94% sensitivity and 95% specificity, compared with a reference standard. Test results were "positive" (presence of *H. pylori* IgG) or "negative" (absence of *H. pylori* IgG).¹⁶ Another portion of the stool sample was analyzed for parasitic infection, utilizing the direct smear method with Lugol's solution and eosin staining, and the Kato-Katz concentration technique for identifying and quantifying helminth eggs.¹⁷

Data analysis

The information was collected in a database and obvious errors and non-plausible data were cleaned. The statistical analysis was performed using Stata v12 (StataCorp, Texas, USA) software and survey commands to allow the use of a planned sample design. Descriptive statistics were calculated (absolute frequencies, percentages) and a bivariate analysis was done using 2×2 contingency tables. The prevalence of infection and its 95% confidence interval were calculated in those tables for each exposure category of the variables, interpreting the statistical association when the interval did not contain the unit. The significant variables, and other scientifically plausible ones, were included

Table 1 Description of the study sample.

Variables	Categories	n	%
Sex	Male	668	52
	Female	606	48
Skin tone	Light	570	45
	Medium	539	42
	Dark	165	13
Municipality	Habana del Este	435	34
	Cerro	176	14
	La Lisa	153	12
	Arroyo Naranjo	510	40
<i>Helicobacter pylori</i> test	Positive	70	5
	Negative	1204	95

in the modeling through dichotomous logistic regression. The result of the *H. pylori* test was the dependent variable, the probability that it was positive was modeled, and prevalence ratios and their 95% confidence intervals adjusted for each exposure variable were obtained, calculating robust standard errors adjusted by municipality. Sex and anti-parasitic treatment were considered a priori confounders but were not introduced into the final model because they did not modify the association by 10% or more.

Ethical considerations

Written statements of informed consent were obtained from the parents or guardians of the underage patients for their participation in the study and the study protocol was approved by the scientific committees of the *Instituto Nacional de Higiene, Epidemiología y Microbiología de La Habana* and the University of Nottingham Medical School in the United Kingdom. The authors declare that no personal information that could identify patients is contained herein.

Results

Of the 1,274 three-year-old children enrolled in the study, 100% of their parents or guardians correctly completed the questionnaire applied by the family physicians. The characteristics of the study participants are shown in [Table 1](#). The prevalence of *H. pylori* infection was 5%.

[Table 2](#) shows the bivariate associations with *H. pylori* and [Table 3](#) shows the factors associated with *H. pylori* infection, after the confounding factor adjustment. The risk factor found for *H. pylori* infection was sleeping together (PR 1.27; 95% CI: 1.03-1.50), and the protective factors were married/living with a partner (PR 0.94; 95% CI: 0.85-0.99) and drinking water delivered by a water truck (PR 0.16; 95% CI: 0.04-0.72).

Discussion and conclusions

To the best of our knowledge, the present analysis is the first population-based study investigating the prevalence of and factors associated with *H. pylori* infection in a population of healthy 3-year-old children in the capital city of Havana, Cuba. The prevalence of bacterial infection in the study population was the result of environmental and

Table 2 Bivariate analysis of exposure to and risk for positive *H. pylori* Ag.

Variables	Categories	Positive <i>Hp</i> test (n = 70) n (%**)	Negative <i>Hp</i> test (n = 1204) n (%**)	PR (95% CI)	p
Paid working mother	No	32 (7)	458 (93)	1.34 (0.85-2.13)	0.247
	Yes	38 (5)	746 (95)	1	
Mother's educational level	Primary/Secondary	21 (6)	327 (94)	1.46 (0.63-3.36)	0.373
	Pre-university	42 (6)	715 (94)	1.34 (0.61-2.93)	0.461
	University	7 (4)	162 (96)	1	
Mother's civil status	Single	13 (5)	254 (95)	0.53 (0.23-1.20)	0.125
	Married/living with a partner	48 (5)	861 (95)	0.58 (0.29-1.14)	0.122
	Divorced/separated	9 (9)	89 (91)	1	
Family income (NC)	< 225	8 (11)	87 (7)	2.66 (0.90-7.89)	0.067
	225-499	31 (44)	535 (44)	1.73 (0.68-4.38)	0.237
	500-999	26 (37)	429 (36)	1.81 (0.71-4.62)	0.208
	>= 1000	5 (7)	153 (13)	1	
Weight-for- age	< 10 percentile	2 (4)	49 (96)	0.69 (0.17-2.75)	0.845
	10-90 percentile	42 (6)	693 (94)	1	
	> 90 percentile	26 (5)	462 (95)	0.93 (0.58-1.50)	0.773
Height-for-age	< 10 percentile	4 (3)	118 (97)	0.57 (0.21-1.55)	0.529
	10-90 percentile	57 (6)	939 (94)	1	
	> 90 percentile	9 (6)	147 (94)	1.01 (0.51-1.99)	0.262
Weight-for-height	< 10 percentile	3 (3)	104 (97)	0.47 (0.15-1.49)	0.405
	10-90 percentile	38 (6)	595 (94)	1	
	> 90 percentile	29 (5)	505 (95)	0.90 (0.57-1.45)	0.676
Brothers or sisters	No	22 (5)	415 (95)	1	
	Yes	48 (6)	789 (94)	1.13 (0.70-1.86)	0.696
Older brothers or sisters	No	37 (6)	623 (94)	1	
	Yes	33 (5)	581 (95)	0.95 (0.61-1.51)	0.954
Type of family	Functional	57 (5)	1044 (95)	1	
	Dysfunctional	13 (8)	160 (92)	1.45 (0.81-2.59)	0.283
Housing conditions	Good	48 (6)	791 (94)	1	
	Regular	18 (5)	343 (95)	0.87 (0.51-1.48)	0.609
	Poor	4 (5)	70 (95)	0.94 (0.35-2.55)	0.911
Overcrowding	No	45 (5)	794 (95)	1	0.877
	Yes	25 (6)	410 (94)	1.07 (0.67-1.72)	

Table 2 (Continued)

Variables	Categories	Positive <i>Hp</i> test (n = 70) n (%**)	Negative <i>Hp</i> test (n = 1204) n (%**)	PR (95% CI)	p
Children's circle or caretaker's home attendance	No	22 (6)	349 (94)	1	0.763
	Yes	48 (5)	855 (95)	0.90 (0.55-1.46)	
Sleeps accompanied	No	27 (4)	597 (96)	1	0.095
	Yes	43 (7)	607 (93)	1.53 (1.00-2.42)	
Treatment for other parasites	No	60 (5)	1115 (95)	1	0.062
	Yes	10 (10)	89 (90)	1.98 (1.05-3.74)	
Treatment with antibiotics	No	26 (6)	406 (94)	1	0.647
	Yes	44 (5)	798 (95)	0.87 (0.54-1.39)	
Pets	No	43 (6)	723 (94)	1	0.918
	Yes	27 (5)	481 (95)	0.95 (0.59-1.51)	
Dogs	No	51 (6)	811 (94)	1	0.410
	Yes	19 (5)	393 (95)	0.78 (0.47-1.30)	
Cats	No	66 (5)	1144 (95)	1	1.000
	Yes	4 (6)	60 (94)	1.15 (0.43-3.05)	
Other animals	No	59 (5)	1080 (95)	1	0.218
	Yes	11 (8)	124 (92)	1.57 (0.85-2.92)	
Rodents	No	55 (6)	926 (94)	1	0.861
	Yes	15 (5)	278 (95)	0.95 (0.52-1.59)	
Vectors	No	47 (5)	831 (95)	1	0.844
	Yes	23 (6)	373 (94)	1.08 (0.67-1.76)	
Type of water consumed	Bottled	4 (12)	29 (88)	1	0.124
	Aqueduct	55 (6)	910 (94)	0.47 (0.17-1.26)	
	Well	10 (4)	216 (96)	0.37 (0.13-0.97)	
	Tanker	1 (2)	49 (98)	0.17 (0.03-0.69)	
Boils water	No	8 (3)	224 (97)	1.73 (0.84-3.55)	0.176
	Yes	62 (6)	980 (94)	1	

Table 2 (Continued)

Variables	Categories	Positive <i>Hp</i> test (n = 70) n (%**)	Negative <i>Hp</i> test (n = 1204) n (%**)	PR (95% CI)	p
<i>Child's habits</i>					
Plays with soil or sand	No	40 (5)	797 (95)	1	0.155
	Yes	30 (7)	407 (93)	1.43 (0.91-2.27)	
Eats soil or sand	No	67 (5)	1177 (95)	1	0.490
	Yes	3 (10)	27 (90)	1.86 (0.62-5.57)	
Sucks his/her thumb	No	56 (5)	1028 (95)	1	0.291
	Yes	14 (7)	176 (93)	1.43 (0.81-2.51)	
Bites/eats fingernails	No	49 (5)	873 (95)	1	0.750
	Yes	21 (6)	331 (94)	1.12 (0.68-1.84)	
Walks barefoot on the ground	No	48 (5)	890 (95)	1	0.400
	Yes	22 (7)	314 (93)	1.28 (0.76-2.09)	
Scratches his/her anus	No	49 (5)	910 (95)	1	0.363
	Yes	21 (7)	294 (93)	1.30 (0.80-2.14)	
Washes hands after urinating/defecating	No	18 (7)	252 (93)	1	0.423
	Yes	52 (5)	952 (95)	0.78 (0.46-1.31)	
Washes hands before eating	No	7 (5)	130 (95)	1	0.991
	Yes	63 (6)	1074 (94)	1.08 (0.51-2.32)	
Washes hands after playing with soil or sand	No	17 (5)	296 (95)	1	1.000
	Yes	53 (6)	908 (94)	1.02 (0.60-1.73)	
<i>Other parasitic infections</i>					
Parasitic infection	No	5 (14)	1172 (86)	1	0.071
	Yes	65 (5)	32 (95)	2.57 (1.10-6.01)	
Helminthiasis	No	69 (5)	1200 (95)	1	0.658
	Yes	1 (20)	4 (80)	3.68 (0.63-21.6)	
Protozoa	No	4 (13)	1176 (87)	1	0.171
	Yes	66 (5)	28 (95)	2.35 (0.91-6.06)	

95% CI: 95% confidence interval; PR : Prevalence ratio.

%*: Percentage with respect to the total number of children (n = 1274).

%**: Percentage with respect to the total per exposure category.

The p values were obtained from association tests based on the X² distribution.

Table 3 Multivariate analysis of exposure to and risk for positive *H. pylori* Ag.

Variables	Categories	Positive <i>Helicobacter pylori</i> test (n=70) n (%**)	OR (95% CI) for positive <i>Helicobacter pylori</i> test	p
Sleeps accompanied	No	27 (38.6)	1	0.027
	Yes	43 (61.4)	1.27 (1.03-1.50)	
Mother's civil status	Single	13 (18.5)	0.93 (0.80-1.50)	0.061
	Married/living with a partner	48 (68.6)	0.94 (0.85-0.99)	
	Divorced	9 (12.9)	1	
Type of water consumed	Bottled	4 (5.7)	1	0.002
	Aqueduct	55 (78.6)	0.48 (0.16-1.33)	
	Well	10 (14.3)	0.41 (0.12-1.28)	
	Tanker	1 (1.4)	0.16 (0.04-0.72)	

The p values were obtained from Wald significance tests based on χ^2 distribution.

$$\text{Definitive model: } P(Y = 1) = \frac{1}{1 + e^{-\left(\begin{array}{l} -3.1465 + 0.2342 * \text{SleepsAccompanied} \\ -0.2492 * \text{SingleMother} - 0.1776 * \text{MarriedMother} \\ +0.1164 * \text{AqueductWater} \\ -0.0535 * \text{WellWater} - 0.9912 * \text{TankerWater} \end{array} \right)}}$$

lifestyle conditions. The cross-sectional analysis conducted on the 3-year-old children appearing in a database of the longitudinal *HINASIC* cohort provided the knowledge of the magnitude of the infection at that age. Sleeping accompanied was an important risk factor and the civil status of married/living with a partner and drinking water from a water-delivery truck were protective factors.

From the total number of 3-year-old children surveyed, 83% provided the stool samples needed for *H. pylori* testing, which enabled the prevalence of the infection to be calculated. The strengths of our study were the fact that the questionnaires were correctly formulated and applied by qualified medical personnel that had been following the children for several years, the majority of questionnaires were answered by the children's parents (90%), and an internationally validated *H. pylori* test recommended for epidemiologic studies on that age group was employed.¹⁸ In addition, highly qualified technical personnel from the *Instituto Nacional de Higiene, Epidemiología y Microbiología de Cuba (INHEM)* carried out the external control of the results of the parasitology tests performed at the parasitology laboratories of the health areas involved, guaranteeing the reliability of the stool test results. The following two limitations of our study were identified: a history of ulcers or gastritis in the parents or guardians of the participating children was not determined and thus could not be analyzed as an associated factor, and the low prevalence of *H. pylori* Ag made it difficult to analyze the possible associated factors. Nevertheless, the results are valuable.

Cuba is a country that has medium and low resource settings. Despite those conditions and the strong economic blockade imposed by the United States for more than 50 years, unlike other countries with the same level of development, its health and education indicators are similar to those of the developed world. That situation, the fact that our study was conducted in an urban zone, and the high rate of antibiotic use in the pediatric population most likely were contributing factors to our study results.¹⁹⁻²² The prevalence

of *H. pylori* in the 3-year-old children analyzed places us in an intermediate prevalence pattern in the international context, but we are close to the pattern of industrialized countries, if we take into account that early childhood prevalence is approximately 1.2% in Western Europe.⁶ In the Latin American countries of Ecuador, Chile, Brazil, and Mexico, an important percentage of children are already infected (63%, 25%, 31.1% and 10%, respectively).^{21,23-26} On the African continent, studies by Amberbir et al. describe a prevalence of 41% in children 3 years of age in an Ethiopian population, and in Uganda, 28.7% in 1-year-old children, 46% of children between 1 and 2 years of age, and 51.7% of children between 3 and 5 years old are already infected.^{27,28} In a recent meta-analysis conducted by Zabala Torres et al., those authors report prevalence of 20% in children under 6 years of age (95% CI: 14%-25%), among other results.²⁹

Children sleeping together with other persons was the risk factor identified in the present study. Regardless of not having determined a history of infection in the adults living with the children, a high prevalence of infection detected in the adult Cuban population (75-90%)^{8-12,14} and evidence of infection transmission between couples suggest an increased risk for transmission to children living with infected individuals.^{24,25,29-32} Family has been found to play a fundamental role in the transmission of the *H. pylori* bacterium, especially within the first years of life, during which the most probable cause is oral-oral transmission from the mother to the child.^{24,29,32}

It appears that living in a stable nuclear family unit is a protective factor against bacterial infection. Greater care given to the children is most likely an important factor and a plausible explanation is the presence of better personal and environmental hygiene habits related to a higher educational level of the study population. In addition, the greater antibiotic use detected in the study participants was relevant, which could lend important weight to that result.^{24,31}

The consumption of potable water delivered by water truck (tankers) was another factor that contributed to the

protection against *H. pylori* infection. The technical condition of the water supply system is not good in all parts of Havana and therefore the distribution of water by tankers is necessary. Despite the inconvenience of that type of public service, it appears that water is safer for drinking, if it does not flow through the water pipeline network, where it can become cross-contaminated due to structural deficiencies. In their 1997 study, Johnson et al. showed that chlorinated water was a factor that inactivated the bacterium, interfering with its transmission by water.³³ A recent review revalidated contaminated water as a source of *H. pylori*, but insists on the need for further detailed study on that route of transmission.³⁴ Eichelberger et al. presented information on the association between environmental risk factors and *H. pylori* infection in the United States based on data from the National Health and Nutrition Examination Survey (NHANES), indicating that environmental exposure (incorrect water use and occupational contact with soil) play an important role in the transmission of *H. pylori* (adjusted odds ratio [OR] 2.7, 95% CI: 1.3-5.6).³⁵

In conclusion, the estimated prevalence of *H. pylori* infection in early childhood places Havana in an intermediate position in the international context. Socioeconomic conditions appear to be important factors in the presence of the infection. Opportune interventions should be implemented to reduce future complications.

Financial disclosure

The present study was funded by the Wellcome Trust (090375), the Nottingham University Hospital Charitable Trust, the Nottingham Respiratory Biomedical Research Unit, and the *Instituto Nacional de Higiene, Epidemiología y Microbiología*, Havana, Cuba.

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgements

The authors wish to thank the Natural History of Wheezing in Cuba (*HINASIC*, the Spanish acronym) study group, made up of the following members:

Instituto Nacional de Higiene, Epidemiología y Microbiología

Silvia J Venero-Fernández, Ramón Suárez-Medina, Hermes Fundora-Hernández, Lenina Menocal-Heredia, Yuria Isabel Caraballos-Sánchez, Reina Amelia Quintana, Félix Manuel Rosado-García, Patricia Varona-Pérez, María del Carmen Hinojosa.

Hospital Universitario Pediátrico Docente Centro Habana Regla N. Rivero, Javier Muñoz-Pérez, Caridad González-Morfa.

Municipio de Arroyo Naranjo

Esperanza de la C. Mora-Faife, Damaris Zaldívar-Ricardo, Maritza Diburt-Amíta, Gisela Álvarez-Valdez, Anadelis Alfonso-Hernández, Vilma Álvarez-Valdez, Yamilka Magaña-Álvarez, Zoe de los Ángeles Figueroa-Barreto, Nieves Sardiñas-Báez, Jorge Antonio Febles-del-Toro, Yunia Velázquez-Pérez, Manuel Hugo Felpeto-Fuentes, Yordanka

Gainza-Bueno, Grisel M. Esquivel-Barrios, Maite Suárez-Paz, Bárbara Judith Magaña-Álvarez, Alida Carménate-Fernández, Roberto Hidalgo-Mederos, Lianet del Loreto, Hidalgo Mederos, Danay Silva, Gretel Comas Fonseca, Dalia M Lazaga Cala, Cristina Odalys, Kessel Díaz, Aliuska Lorenzo Méndez

Municipio de La Lisa

Gladys García-García, María de Lourdes Ortiz-Hernández, María Antonia Betancourt-López, Marlén Batista-Cedeño, Iris Alfonso-Castellanos, Leticia Gómez-García, Ernesto Rafael Gutiérrez-Mendoza, María Luisa Loynaz-González, Nibenia Rodríguez-Trujillo, Yanet Pozo-Herrera, Víctor Manuel Montejo-Guerra, Julia Urbina-Reynaldo, Valentina Gómez-Suliman, Caridad Alicia Rodríguez-Aragón.

Municipio de Cerro

Ileana del Valle-Infante, Claudia Matos-Ramos, Martha Betancourt-Orue, Oscar Alba-Monteagudo, Yuderkis Ferrer-Ceruto, Aída Damas-Martínez, Mercedes Peñalver-Pérez

Municipio de Habana del Este

Liem Gómez-Marrero, Sarahí Castillo-Martínez, Amor de los Angeles Castaño-Vega, Norberto Torriente-Barzaga, Ileana Ávila-Rodríguez, Magalys Navarro-Ruiz, Kirenia Díaz-Hernández, Iluska De La Torre Suarez, Gilberto Roque-Pereira, Yamilet Corona-Carnero, Iдания Gonzalaz-Fernandez, Fidelia Romeo-Ravelo, Regla Hernandez-Ponce, Teresa Serrano-González, Dulce Romeo-Cepero, Caridad Gonzalez Leiva, Teresa De Jesus Cobas Espino, Nuris-Fajardo, Midiala Perez-Arcia, Sarahy Diaz-Araujo, Yanet Medina- Lescay, Sandra Collazo-Rodriguez, Julia Amparo Griñán-Ramos, Teresa Serrano-Gonzalez, Beatriz Lazo-Vazquez, Tania Pupo-Portal, Nidia Leyva-Porra, Odalys Pacheco-Mesa, Martha Rizo-Ramos, Yaneysi Vallafuerte-Perez, Aliniuska De La Paz-Arias, Maite B Garcia-Sotolongo, Yusimí Calzado-Herrera, Martha Nidia Rizo-Ramos, Guillermo Verdecia, Mayté B García-Sotolongo, Juana F Abreu-Quijano, Fidelia Romeo-Ravelo.

The authors also wish to thank the children and parents for providing their data and the municipal Public Health directors and laboratory personnel that supported the study.

References

1. Suerbaum S, Michetti P. *Helicobacter pylori* infection. *N Engl J Med*. 2002;347:1175-86. Available from: <http://extranet.who.int/hinari/es/journals.php>
2. McColl KE. Clinical practice. *Helicobacter pylori* infection. *N Engl J Med*. 2010;362:1597-604. Available from: <http://www.nejm.org/doi/pdf/10.1056/NEJMcp1001110>
3. Zullo A, Ridola L, Hassan C, et al. Glaucoma and *Helicobacter pylori*: eyes wide shut? *Dig Liver Dis*. 2012;44:627-8. Available from: <http://extranet.who.int/hinari/es/journals.php>
4. Linz B, Balloux F, Moodley Y, et al. An African origin for the intimate association between humans and *Helicobacter pylori*. *Nature*. 2007;445:915-8. Available from: <http://extranet.who.int/hinari/es/journals.php>
5. Marshall B, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet*. 1984;232:1311-5. Available from: <http://extranet.who.int/hinari/es/journals.php>
6. Crowe SE. Bacteriology and epidemiology of *Helicobacter pylori* infection; 2013. Available from: UpToDate, Inc. UpTo

- Date <http://www.uptodate.com/contents/bacteriology-and-epidemiology-of-helicobacter-pylori-infection>
7. Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. *Nature reviews. Genetics*. 2012;13:260–70 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3418802/>
 8. Muhammad JS, Zaidi SF, Sugiyama T. Epidemiological ins and outs of *Helicobacter pylori*: a review. *J Pak Med Assoc*. 2012;62:955–9. Available from: <http://extranet.who.int/hinari/es/journals.php>
 9. Pascual MGC, Rojas-Zurita F, Grá OB, et al. Prevalencia de la infección por *Helicobacter pylori* en pacientes dispepticos. *Rev Panam Infectol*. 2004;6:8–14. Available from: ID: lil-400901.
 10. Ortiz-Prinz D, Guariglia-orozepe V, Ávila M, et al. *Helicobacter pylori* cagA and vacA genotypes in Cuban and Venezuelan populations. *Mem Inst Oswaldo Cruz*. 2010;105:331–5, <http://dx.doi.org/10.1590/s0074-02762010000300016>. Available from: <http://www.scielo.br/pdf/mioc/v105n3/07.pdf>
 11. Martínez-Echavarría MT, Noa-Pedroso G. Infección por *Helicobacter pylori* en pacientes con mucosa sana y con gastritis erosiva. *Revista Cubana de Medicina*. 2009;48:0-0. Available from: http://bvs.sld.cu/revistas/med/vol48_2_09/medsu209.htm
 12. Ledesma Z, Gutiérrez B, Cirión GR, et al. Diagnóstico histológico de la infección por *Helicobacter pylori* en Pinar del Río, Cuba. *Vacci Monitor*. 2010;19:1–4. Available from: www.finlay.sld.cu/publicaciones/vaccimonitor/Vm2010/a6.pdf
 13. Ruiz Álvarez V, Reboso-Pérez J, Hernández-Triana M. Asociación entre la infección por *Helicobacter pylori* y anemia en niños de edad escolar. *Rev Cuba Invest Biomed*. 2005;24:0-0. Available from: bvs.sld.cu/revistas/ibi/vol24_2_05/ibi02205.pdf
 14. Gámez Escalona MM, Mulet Pérez CAM, Miranda Moles Z, et al. Gastritis crónica antral por *Helicobacter pylori* en la infancia. *Rev Cub.Pediatría*. 2008;80:0-0. Available from: http://bvs.sld.cu/revistas/ped/vol80_1_08/ped02108.htm
 15. Venero Fernandez S, Suarez Medina R, Mora Faife E, et al. Risk factors for wheezing in infants born in Cuba. *Q J M*. 2013;106:1023–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23824939>
 16. Spinreact. Disponible en: http://www.spinreact.com/files/Inserts/Placas/OSIS11_Ref._1504040_H._pylori_ANTIGENS_rev2011.pdf.
 17. Núñez FA, Cordovi Prado R. *Manual de técnicas básicas para el diagnóstico de las parasitosis intestinales*. La Habana: Ministerio de Salud Pública de Cuba/UNICEF; 2004.
 18. Koletzko S, Feydt-Schmidt A. Infants differ from teenagers: use of noninvasive tests for detection of *Helicobacter pylori* infection in children. *Eur J Gastroenterol Hepatol*. 2001;13:1047–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11564953>
 19. Organización Mundial de la Salud, Estadísticas Sanitarias Mundiales. OMS; 203. Available from: <http://www.who.int/gho/publications/world.health.statistics/2013/es/>.
 20. Organización de las Naciones Unidas para la Educación, la Ciencia y la Cultura (UNESCO). *The state of education in Latin America and the Caribbean guaranteeing quality education for all 2010*. Estadística. Paris: UNESCO 2010-2015. Available from: http://www.unicef.org/2fmedia%2Ffiles%2FState_of_education_in_Latin_America_and_Caribbean.pdf&ei=0-5PU8fZBsyW0gHR_IG4Cw&usg=AFQjCNEdguxWUIV6sNNnEs5GcXa245k8Hg&bvwm=bv.64764171.d.dmQ.
 21. O’Ryan ML, Rabello M, Cortés H, et al. Dynamics of *Helicobacter pylori* detection in stools during the first 5 years of life in Chile, a rapidly developing country. *Pediatr Infect Dis J*. 2013;32:99–103, <http://dx.doi.org/10.1097/INF.0b013e318278b929>. Available from: .
 22. Lara Bastanzuri MC, Cires Pujol M, García Miliam AJ. Consumo de antimicrobianos en APS. *Rev Cubana Med Gen Integr*. 2003;19:0-0. Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0864-21252003000400003&lng=es
 23. Gómez NA, Salvador A, Vargas PE, et al. Seroprevalencia de *Helicobacter pylori* en la población infantil ecuatoriana. *Rev Gastroenterol Peru*. 2004;24:230–3. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/?term=G%C3%B3mez+NA%2C+Salvador+A%2C+Vargas+PE%2C>.
 24. da Silva Roque JR, Strehl Machado R, Rodrigues D, et al. Prevalence of *Helicobacter pylori* infection in an indigenous community in São Paulo and associated factors: cross-sectional study. *Sao Paulo Med. J*. 2017;135:140–5. Available from: http://www.scielo.br/scielo.php?pid=S1516-31802017000200140&script=sci_arttext
 25. Cartagenes VD, Martins LC, Carneiro LM, et al. *Helicobacter pylori* em crianças e associação de cepas CagA na transmissão mãe-filho na Amazônia brasileira. *Rev Soc Bras Med Trop*. 2009;42:298–302. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0037-86822009000300011&lng=en&nrm=iso&lng=en
 26. Madrazo-de la Garza JA, González-Ortiz B. *Helicobacter pylori* en niños. *Bol Med Hosp Infant Mex*. 2001;58:656–62. Available from: http://www.imbiomed.com.mx/1/1/articulos.php?articulos.php?method=showDetail&id_articulo=3919&id_seccion=362&id_ejemplar=442&id_revista=20
 27. Amberbir A, Medhin G, Erku W, et al. Effects of *Helicobacter pylori*, geohelminth infection and selected commensal bacteria on the risk of allergic disease and sensitization in 3-year-old Ethiopian children. *Clin Exp Allergy*. 2011;41:1422–30, <http://dx.doi.org/10.1111/j.1365-2222.2011.03831.x>. Available from: .
 28. Hestvik E, Tylleskar T, Kaddu-Mulindwa DH, et al. *Helicobacter pylori* in apparently healthy children aged 0-12 years in urban Kampala, Uganda: a community-based cross sectional survey. *BMC Gastroenterology*. 2010;10:62. Available from: <https://bmcgastroenterol.biomedcentral.com/articles/10.1186/1471-230X-10-62>
 29. Zabala-Torres B, Lucero Y, Lagomarcino AJ, et al. Review. Prevalence and dynamics of *Helicobacter pylori* infection during childhood. *Helicobacter*. 2017;22:1–18, <http://dx.doi.org/10.1111/hel.12399>. Available from: .
 30. Rodríguez E, Bermúdez L, Trujillo M, et al. Evidencia de la transmisión de la infección de *Helicobacter pylori* entre parejas. *Revista CENIC. Ciencias Biológicas*. 2010;41:0-0. Available from: <http://revista.cnic.edu.cu/revistaCB/sites/default/files/articulos/CB-2010-4-CB-018.pdf>
 31. Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. *Nature reviews. Genetics*. 2012;13:260–70 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3418802/>
 32. Weyermann M, Rothenbacher D, Brenner H. Acquisition of *Helicobacter pylori* infection in early childhood: independent contributions of infected mothers, fathers, and siblings. *Am J Gastroenterology*. 2009;104:182–9.
 33. Johnson CH, Rice EW, Reasoner DJ. Inactivation of *Helicobacter pylori* by chlorination. *Appl Environ Microbiol*. 1997;63:4969–79. Available from: <http://europepmc.org/backend/ptpmrender.fcgi?accid=PMC168826&blobtype=pdf>
 34. Aziz RK, Khalifa M, Sharaf RR. Contaminated water as a source of *Helicobacter pylori* infection: a review. *J Adv Res*. 2015;6:539–47. Available from: <http://www.sciencedirect.com/science/article/pii/S2090123213000970>
 35. Eichelberger L, Murphy G, Etemadi A, et al. Risk of gastric cancer by water source: evidence from the Golestan case-control study. *PLOS ONE*. 2015;10:e0128491. Available from: <http://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0128491&type=printable>