



REVISTA DE GASTROENTEROLOGÍA DE MÉXICO

www.elsevier.es/rgmx



ORIGINAL ARTICLE

A comparison of endoscopic procedures performed at a tertiary care hospital before and during the SARS-CoV-2 pandemic

F.E. Zárate-Mondragón^{a,*}, M. Esparza-Aguilar^b, J.F. Cadena-León^a,
K.R. Ignorosa-Arellano^a, E. Montijo-Barrios^a, E.M. Toro-Monjaráz^a,
R. Cervantes-Bustamante^a, A.G. Ayala-Germán^a, J.A. Ramírez-Mayans^a

^a Servicio de Gastroenterología y Nutrición, Instituto Nacional de Pediatría, Mexico City, Mexico

^b Servicio de Epidemiología, Instituto Nacional de Pediatría, Mexico City, Mexico

Received 1 August 2023; accepted 24 October 2023

KEYWORDS

COVID pandemic;
Upper and lower
endoscopy;
Pediatrics

Abstract

Introduction and aim: SARS-CoV-2 emerged in 2019 and had a huge impact on the world. The area of endoscopy suffered great changes, causing a reduction in the number of procedures and its indications. The aim of our study was to compare the quantity, indication, and type of procedures in 2019 with those in 2020.

Method: A retrospective, observational, analytic, and cross-sectional study was conducted, obtaining information from the endoscopy registry. The STROBE checklist was employed.

Statistical analysis: The quantitative variables were analyzed with descriptive statistics (measures of central tendency and dispersion) and the categorical variables with frequencies and percentages. The quantitative variables were compared, using the Student's t test/Mann-Whitney U test, and the categorical variables with contingency tables, using the Fisher's exact test.

Results: In 2019, a total of 277 procedures were performed, compared with 139 in 2020. Mean patient age was 98.53 months (61.46 SD) in 2019 and 77.02 months (59.81 SD) in 2020; 352 diagnostic procedures and 136 therapeutic procedures were carried out in 2019, compared with 51 diagnostic procedures and 88 therapeutic procedures in 2020. The number of diagnostic and therapeutic procedures were inverted (72.1%–36.7% and 27.9%–63.3%, respectively) ($p < 0.0001$). Esophageal varices, upper gastrointestinal bleeding (UGIB), and foreign body extraction were the indications, in order of predominance in 2019, compared with foreign body extraction ($p < 0.05$), UGIB, and esophageal varices in 2020. There were no differences regarding colonoscopy.

* Corresponding author at: Insurgentes Sur 3700-C, Insurgentes Cuicuilco, cp 04530. Tel.: 5535120580.

E-mail address: florazarate@gmail.com (F.E. Zárate-Mondragón).

PALABRAS CLAVE

Pandemia COVID;
Endoscopia alta y
baja;
Pediatria

Conclusion: There was a clear difference in indication and type of procedure, with an increase in foreign body extraction in preschoolers.

© 2024 Published by Masson Doyma México S.A. on behalf of Asociación Mexicana de Gastroenterología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Comparación en los procedimientos endoscópicos antes y durante la pandemia del SARS-CoV-2 en un tercer nivel de atención

Resumen

Introducción: En 2019 surgió el SARS-CoV-2 que tuvo un gran impacto a nivel mundial, el área de endoscopia sufrió grandes cambios provocando una reducción del número de procedimientos y sus indicaciones.

Objetivo: Comparar: cantidad, indicación y tipo de procedimientos del 2019 contra 2020.

Método: Estudio observacional, analítico, transversal y retrospectivo obteniendo la información del registro de endoscopia. Se utilizó la lista de cotejo STROBE.

Análisis estadístico: Variables cuantitativas analizadas con estadística descriptiva (medidas de tendencia central y dispersión) y para las categóricas, frecuencias y porcentajes. Para comparar se utilizó T de Student/U de Mann-Whitney para las variables cuantitativas; tablas de contingencia con prueba Ji cuadrada o exacta de Fisher para categóricas.

Resultados: En el 2019 se realizaron 277 procedimientos en comparación de 139 en 2020. Media de edad en el 2019 fue de 98.53 meses (DE 61.46) y para el 2020 la media fue 77.02 (DE 59.81), tipo de procedimiento en 2019, 352 fueron procedimientos diagnósticos y 136 terapéuticos, mientras que, en 2020, 51 fueron diagnósticos y 88 terapéuticos. Se invirtió la proporción de procedimientos diagnósticos (72.1% al 36.7%) y terapéuticos (27.9% a 63.3%), ($p < 0.0001$). Las indicaciones en 2019 predominaron vórices esofágicas, hemorragia de tubo digestivo alto (HTDA) y extracción de cuerpo extraño, en comparación con el 2020 donde predominó extracción de cuerpo extraño ($p < 0.05$), HTDA y vórices esofágicas. En la colonoscopia no hubo diferencias.

Conclusión: Hubo una clara diferencia en la indicación y tipo de procedimiento, hubo un incremento en la extracción de cuerpos extraños en pre-escolares.

© 2024 Publicado por Masson Doyma México S.A. en nombre de Asociación Mexicana de Gastroenterología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In 2019, a new highly contagious virus named SARS-CoV-2, associated with severe acute respiratory syndrome, emerged.¹⁻³ In January 2020, the World Health Organization (WHO) declared the presence of this virus to be a Public Health Emergency of International Concern, and in March of the same year, a pandemic.² The SARS-CoV-2 world crisis has negatively impacted the number and type of surgical procedures performed due to the need to recategorize emergency, urgent, and elective procedures.⁴⁻⁶ As a consequence, this brought about an 81.4% reduction in endoscopic procedures, according to a survey conducted by the European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN).⁷

By January 25, 2021, there were an estimated 1,771,749 persons infected with COVID-19 and 150,273 deaths from the disease in Mexico, signifying a lethality of 6%.⁸ This infection rate was reflected at the *Instituto Nacional de Pediatría*, resulting in the temporary closure of the endoscopy units due to a lack of negative pressure and laminar flow. Consequently, procedures have had to be restricted to operating

rooms, and because they must also be used by the rest of the surgical services, this has caused an important decrease in the performance of endoscopic procedures.

The aim of the present study was to compare the number and type of endoscopic procedures performed at the Gastroenterology and Nutrition Service of the *Instituto Nacional de Pediatría*, before and during the SARS-CoV-2 pandemic.

Materials and methods

A retrospective, observational, analytic, and cross-sectional study was conducted at the *Instituto Nacional de Pediatría*. The STROBE checklist for cross-sectional studies was utilized. Records from January to December of 2019 from the area of endoscopy were reviewed and compared with the records from 2020. The demographic characteristics, indications for the procedure, type of procedure, and endoscopic findings were obtained.

Because of its retrospective design, the study was classified as negligible-risk research. Based on the Regulation of the General Health Law in Health Research, Second Title, of




Figure 1 Monthly frequency of endoscopic procedures between 2019 and 2020 in patients at the Instituto Nacional de Pediatría.

the Ethical Aspects in Research on Humans, Chapter 1, Article 23, given that this is a negligible-risk analysis, informed consent was not requested.

Statistical analysis

The analysis was carried out, using the SPSS version 22.0 program. Through descriptive statistics, the quantitative variables were expressed as measures of central tendency and dispersion and the categorical variables as frequencies and percentages. The Student's t test or Mann-Whitney test were utilized to compare the quantitative variables, and contingency tables with the Fisher's exact test to compare the categorical variables.

Ethical considerations

The study was conducted following the international regulations in bioethics and met the norms of the *Instituto Nacional de Pediatría*. It was reviewed by the institutional ethics committee. No experiments were carried out on humans or animals and there was complete data confidentiality. This study is classified as a negligible-risk study, given that clinical records were reviewed, retrospective documental research methods and techniques were employed, and no interventions on or intentional modifications of the physiologic, psychologic, and social variables of the individuals that participated in the study were performed, thus informed consent was not required.

Results

Upper endoscopy

In the descriptive analysis, a total of 277 upper endoscopy procedures were performed in 2019, compared with 139 procedures in 2020 (Fig. 1). In 2019, mean patient age was 98.53 months (61.46 standard deviation [SD]), whereas in 2020, it was 77.02 months (59.81 SD). In 2019, the procedures were performed on 277 males and 213 females, whereas they were performed on 79 males and 60 females in 2020. In 2020, there was no statistically significant difference by sex, with respect to the number of patients treated ($p > 0.999$).

Table 1 Comparison of patients divided into age groups.

	Fisher's exact test		
	2019	2020	p value
Neonates	0.4%	0.0%	>0.9999
Infants	19.0%	26.6%	0.0575
Preschoolers	20.2%	34.5%	0.0006
School-age children	36.7%	19.4%	<0.0001
Adolescents	23.7%	19.4%	0.3051

The patients were divided into age groups and Table 1 shows the comparison made between 2019 and 2020, using the Fisher's exact test.

Regarding the type of procedure, in 2019, 352 were diagnostic procedures and 136 were therapeutic, whereas in 2020, 51 were diagnostic and 88 were therapeutic, resulting in an inversion in the number of diagnostic procedures (from 72.1% to 36.7%) and therapeutic procedures (from 27.9% to 63.3%), ($p < 0.0001$).

With respect to indications for upper endoscopy, in 2019, the order of predominance was esophageal varices, upper gastrointestinal bleeding (UGIB), and foreign body extraction, compared with foreign body extraction, UGIB, and esophageal varices in 2020. Table 2 shows all the indications.

Table 3 shows that esophagitis and malformations were the findings that were significantly increased in 2020.

There was a significant decrease in 2020, in the number of patients in whom no procedures were performed but biopsies were taken. There was also a decrease in banding ligation/sclerotherapy, a slight increase in hemostasis, and a considerable increase in foreign body extraction in 2020, all with statistical significance (Table 4).

The median number of upper endoscopies in 2020 was 9.5 procedures per month (minimum: 2, maximum: 23), which was significantly lower than the 43 procedures per month in 2019 (minimum: 12, maximum: 56) ($p = 0.0001$ in the Mann-Whitney U test).

Colonoscopy

In the two years analyzed, 150 procedures were performed; 104 in 2019 and 46 in 2020, resulting in a decrease of 66%. Regarding demographic characteristics, in 2019, the procedures were carried out on 51 males and 53 females, whereas in 2020, they were carried out on 28 males and 18 females. There was no difference by sex between the two years ($p = 0.02160$). Mean patient age was 109.61 months (58.26 SD) in 2019, whereas it was 109.20 months (64.95 SD) in 2020. The patients were divided by age group, which highlighted the fact that preschoolers were more affected in 2020, compared with school-age children, in whom there was a significant reduction. The findings are shown in Table 5.

In 2019, 76 diagnostic colonoscopies and 28 therapeutic colonoscopies were performed, whereas 29 diagnostic colonoscopies and 17 therapeutic colonoscopies were performed in 2020, signifying a decrease of 62% in the diagnostic procedures and 40% in therapeutic ones. No proportional

Table 2 Indications for upper endoscopy.

Clinical presentation	Fisher's exact test		p value
	2019 n (%)	2020 n (%)	
GER/vomiting	41 (8.4%)	3 (2.2%)	0.0081
Dysphagia	25 (5.1%)	1 (0.7%)	0.0161
Achalasia	5 (1%)	2 (1.4%)	0.6533
Esophageal stricture	37 (7.6%)	16 (11.5%)	0.1650
Altered swallowing mechanics	41 (8.4%)	11 (7.9%)	>0.9999
Esophageal varices	92 (18.8%)	20 (14.4%)	0.2597
UGIB	65 (13.3%)	12 (8.6%)	0.1861
Foreign body	55 (11.2%)	48 (34.5%)	<0.0001
Caustic substance intake	14 (2.9%)	3 (2.2%)	>0.9999
CAP	43 (8.8%)	3 (2.2%)	0.0139
Chronic diarrhea	6 (1.2%)	5 (3.6%)	0.0717
IBD	5 (1%)	1 (0.7%)	>0.9999
Probable celiac disease	13 (2.7%)	0 (0%)	0.0829
Polypoid syndromes	13 (2.7%)	0 (0%)	0.0829
LGIB	2 (0.4%)	0 (0%)	>0.9999
Obstructive cause	13 (2.7%)	6 (4.3%)	0.3966
Others	8 (1.6%)	2 (1.4%)	>0.9999
Esophagitis/ulcers/peptic acid disease	8 (1.6%)	2 (1.4%)	>0.9999
Gastrostomy/transpyloric catheter placement	9 (1.8%)	1 (0.7%)	0.6998
Total	490	139	

CAP: chronic abdominal pain; GER: gastroesophageal reflux; IBD: inflammatory bowel disease; LGIB: lower gastrointestinal bleeding; UGIB: upper gastrointestinal bleeding.

Table 3 Endoscopic findings.

Finding	Fisher's exact test		p value
	2019 n (%)	2020 n (%)	
Normal	91 (18.6%)	22 (15.8%)	0.5317
Esophagitis	38 (7.8%)	34 (24.5%)	<0.0001
Esophagitis due to a caustic agent	9 (1.8%)	2 (1.4%)	>0.9999
Esophagitis due to candida	3 (0.6%)	0 (0%)	>0.9999
Stricture	24 (4.9%)	4 (2.9%)	0.362
Varices/congestive gastropathy	107 (21.8%)	25 (18%)	0.3476
Barrett's esophagus/tumors	5 (1%)	1 (0.7%)	>0.9999
Esophageal dilation	0 (0%)	1 (0.7%)	0.2210
Malformations	3 (0.6%)	5 (3.6%)	0.0155
Achalasia	1 (0.2%)	0 (0%)	>0.9999
Esophageal ulcer	11 (2.2%)	3 (2.2%)	>0.9999
Gastric ulcers	21 (4.3%)	5 (3.6%)	>0.9999
Foreign body	16 (3.3%)	8 (5.8%)	0.2074
Biliary reflux	3 (0.6%)	0 (0%)	0.2658
Variceal sequelae	1 (0.2%)	0 (0%)	>0.9999
Polyps	7 (1.4%)	2 (1.4%)	>0.9999
Nonerosive gastropathy	46 (9.4%)	7 (5%)	0.1201
Erosive gastropathy	38 (7.8%)	7 (5%)	0.3517
Duodenitis	14 (2.9%)	1 (0.7%)	0.2106
Nodular gastropathy	25 (5.1%)	2 (1.4%)	0.0607
Hemorrhagic gastropathy	11 (2.2%)	1 (0.7%)	0.4798
Hypertrophic folds	2 (0.4%)	1 (0.7%)	0.5279
Fundoplication	2 (0.4%)	0 (0%)	>0.9999
Others	12 (2.4%)	8 (5.8%)	0.0582
Total	490	139	

Table 4 Type of procedure.

Procedure	2019 n (%)	2020 n (%)	Fisher's exact test p
None	152 (31%)	30 (21.6%)	0.0339
Banding ligation/sclerotherapy	26 (5.3%)	18 (12.9%)	0.0034
Dilations	30 (6.1%)	13 (9.4%)	0.1855
Hemostasia (clips, hemospray, adrenaline)	5 (1%)	6 (4.3%)	0.0181
Transpyloric catheter placement	3 (0.6%)	3 (2.2%)	0.1253
Gastrostomy placement, change, or removal	32 (6.5%)	12 (8.6%)	0.4502
Polypectomy	5 (1%)	0 (0%)	0.589
Foreign body extraction	28 (5.7%)	34 (24.5%)	<0.0001
Botulinum toxin	2 (0.4%)	1 (0.7%)	0.2604
Biopsy	207 (42.2%)	21 (15.1%)	<0.0001
Total	490	139	

Table 5 Frequency, number, and comparison of colonoscopies by age group.

Age	Fisher's exact test		p value
	2019 n (%)	2020 n (%)	
Infant	4 (3.8%)	3 (6.5%)	0.6923
Preschooler	23 (22.1%)	13 (28.2%)	0.0168
School-age child	46 (44.2%)	15 (32.6%)	0.0403
Adolescent	31 (29.8%)	15 (32.6%)	0.0503
Total	104	46	

differences were detected ($p = 0.2843$), with diagnostic procedures always predominating.

Table 6 shows the indications, diagnoses, and procedures in colonoscopy. No statistically significant differences were detected in the comparison of the two years.

In 2020, only 31% (185) of the procedures were performed, compared with 594 procedures in 2019, implying a decrease of 69%. When dividing the procedures, 139 (28%) upper endoscopies were performed in 2020, compared with 490 in 2019, resulting in a 72% decrease; 104 colonoscopies were performed in 2019 and 46 (44%) in 2020, signifying a 66% decrease.

The median number of colonoscopies in 2020 of 3.5 monthly procedures (minimum: 0, maximum: 15) was significantly lower than the 10 monthly procedures in 2019 (minimum: 1, maximum 15) $p = 0.0199$ in the Mann–Whitney U test) (**Fig. 1**).

Discussion

In the medical literature, the reported impact of the pandemic on the decrease in the number of procedures is on average 80%. In our study, we found that endoscopic procedures were reduced by 69%. Urgent and emergency procedures were predominant during the pandemic, which is a protocol that has been continued by the *Instituto Nacional de Pediatría*, in accordance with recommendations by NASPGHAN, ESPGHAN, and the GESA.^{6,9–11}

When comparing age at the time of endoscopy, school-age children predominated in 2019 and decreased in 2020 ($p < 0.05$), whereas the number of preschoolers was higher in 2020 ($p < 0.05$). This is explained by the longer time chil-

dren remained at home, often unsupervised, accounting for the increase of foreign body ingestion in children. In addition, foreign body extraction is an urgent or emergency procedure, and in the 2019 and 2020 comparison, suspected foreign body was one of the most important indications in 2020 ($p < 0.05$).

One of the differences between 2019 and 2020 is the type of procedure performed. In 2019, diagnostic endoscopy was more frequently performed, whereas in 2020, it was therapeutic endoscopy ($p < 0.05$). Once again, this is explained by the indication to only perform urgent and emergency procedures, resulting in a decrease in diagnostic endoscopy of 86%, a figure coinciding with that reported by the ESPGHAN,¹¹ whereas therapeutic endoscopy decreased by only 36%.

Among the main medical indications for performing upper endoscopy in 2019 were: esophageal varices (19%) (92/490), gastrointestinal bleeding (13%) (65/490), suspected foreign body (11%) (55/490), altered swallowing mechanics (8%) (41/490), gastroesophageal reflux (8%) (41/490), and esophageal stricture (7.5%) (37/490). In 2020, the main indications were: suspected foreign body (34%) (48/139), esophageal varices (14.5%) (20/139), stricture (11.5%) (16/139), UGIB (8.6%) (12/139), altered swallowing mechanics (8%) (11/139), and gastroesophageal reflux (2%) (3/139). In 2020, suspected foreign body was one of the main indications ($p < 0.05$). Notably, variceal and non-variceal bleeding were the indications whose numbers most commonly increased in 2020 ($p < 0.05$), along with stricture.

Likewise, there was an important change in endoscopic findings, given that in 2019, the most common finding was varices/congestive gastropathy (26%) (107/490), followed

Table 6 Colonoscopy indications, diagnosis, and procedure performed.

Indication for colonoscopy	2019 n (%)	2020 n (%)	Fisher's exact test p value
UGIB	35 (33.7%)	20 (43.5%)	0.2738
Polyposis	28 (26.9%)	12 (26.1%)	0.5224
Chronic diarrhea	10 (9.6%)	3 (6.5%)	0.755
IBD	15 (14.4%)	7 (15.2%)	>0.9999
Allergic colitis	3 (2.9%)	0 (0%)	0.5531
Tumors	2 (1.9%)	0 (0%)	>0.9999
CAP	3 (2.9%)	1 (2.2%)	>0.9999
GVHD	4 (3.8%)	2 (4.3%)	>0.9999
Others	2 (1.9%)	1 (2.2%)	>0.9999
Infectious colitis	2 (1.9%)	0 (0%)	>0.9999
Total	104	46	
<i>Colonoscopic diagnosis</i>			
Normal	25 (24%)	6 (13%)	0.1885
Polyps	34 (32.7%)	21 (45.7%)	0.1443
Nonspecific inflammation	44 (42.3%)	14 (30.4%)	0.2045
IBD	0 (0%)	1 (2.2%)	0.3067
Anatomic malformations	0 (0%)	3 (6.5%)	0.0275
AVM	1 (1%)	1 (2.2%)	0.5207
Total	104	46	
<i>Colonoscopic procedure</i>			
None	1 (1%)	0 (0%)	>0.9999
Biopsy	94 (90.4%)	41 (89.1%)	0.7759
Polypectomy	8 (7.7%)	3 (6.5%)	>0.9999
Hemostasis	0 (0%)	1 (2.2%)	0.3067
Botulinum toxin	1 (1%)	1 (2.2%)	0.5207
Total	104	46	

AVM: arteriovenous malformation; CAP: chronic abdominal pain; GVHD: graft-versus-host disease; IBD: inflammatory bowel disease; UGIB: upper gastrointestinal bleeding.

by normal endoscopy (18.5%) (91/490), which is explained by the fact that the indications in that year were mainly diagnostic: nonerosive gastropathy (9%) (46/490), esophagitis (7.7%) (38/490), and erosive gastropathy (7.7%) (38/490), whereas in 2020, the most common findings were esophagitis (24%) (34/139), varices/congestive gastropathy (18%) (25/139), normal endoscopy (16%) (22/139), foreign body (6%) (8/139), and erosive and nonerosive gastropathy, both with (5%) (7/139). There were more abnormal (pathologic) studies than normal studies in 2020, compared with 2019 ($p < 0.05$), again explaining the changes in endoscopy indications in the two years.

There was also an important change in the type of procedure performed. Compared with 2020, diagnostic biopsy (42%) (207/490) was the most frequent procedure in 2019, followed by no procedure (31%) (152/490), dilations (6%) (30/490), foreign body (5.7%) (28/490), and banding ligation/sclerotherapy (5.3%) (26/490). In 2020, the main procedure was foreign body extraction (24%) (34/139), followed by no procedure (21%) (30/139), biopsy (15%) (21/139), banding ligation/sclerotherapy (13%) (18/139), and dilations (9%) (13/139). There was a statistically significant decrease in the number of biopsies due to the fact that in 2020, indications for endoscopy were mainly therapeutic, and not diagnostic. Foreign body extraction ($p < 0.05$) and banding ligation and/or variceal sclerotherapy were the most important procedures in 2020 because children were at home longer, often unsupervised, lead-

ing to a substantial increase in foreign body ingestion, whereas esophageal variceal bleeding continued being an emergency.

Colonoscopy

With respect to colonoscopy, there was a 66% decrease in the procedure in 2020; 73% of the procedures were diagnostic in 2019, dropping to 63% in 2020; therapeutic procedures increased to 37% in 2020, compared with 26% in 2019, as was expected.

The clinical indications were the same for the two years: lower gastrointestinal bleeding (33.6% in 2019 and 43.5% in 2020) and polyposis (27% in 2019 and 26% in 2020), followed by chronic diarrhea, inflammatory bowel disease, and graft-versus-host disease. Even though the proportions were different, the indications remained more or less the same. Bleeding continued to be one of the most common causes both years. No statistically significant differences were found.

In both years, regarding diagnostic colonoscopy, nonspecific inflammation predominated (42% in 2019 and 30% in 2020), followed by polyps (33% in 2019 and 47% in 2020) and normal colonoscopy (24% in 2019 and 13% in 2020). Fewer diagnostic procedures were performed in 2020, thus there were fewer normal studies reported, but those results were not statistically significant.

Conclusions

According to the data reported, there was an important decrease in the number of procedures performed, as well as a change in the type of procedure, with a decrease in diagnostic upper endoscopy and an increase in therapeutic endoscopy.

One of the most common causes for diagnostic and therapeutic upper endoscopies is UGIB and the presence of esophageal varices and hypertensive gastropathy. Clinical indications for colonoscopy remained the same in the two years, with lower gastrointestinal bleeding and polyps being the main causes. This highlights the importance of endoscopy services being supplied with all the new hemostatic techniques (hemospray, hemostasis clips, argon plasma, silver nitrate, etc.)

In addition, due to the lockdown, there was an increase in foreign body ingestion, especially in preschoolers, suggesting the need for reinforcing prevention measures in the home and increasing the awareness of pediatricians of the possibility of this occurrence.

Financial disclosure

No financial support was received in relation to this article.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Patel NA. Pediatric COVID-19: systematic review of the literature. *Am J Otolaryngol.* 2020;41:102573, <http://dx.doi.org/10.1016/j.amjoto.2020.102573>.
2. Budhwar S, Sethi K, Chakraborty M. A rapid advice guideline for the prevention of novel coronavirus through nutritional intervention. *Curr Nutr Rep.* 2020;9:119–28, <http://dx.doi.org/10.1007/s13668-020-00325-1>.
3. Calder PC. Nutrition, immunity and COVID-19. *BMJ Nutr Prev Health.* 2020;3:74–92, <http://dx.doi.org/10.1136/bmjnp-2020-000085>.
4. Chiu PWY, Ng SC, Inoue H, et al. Practice of endoscopy during COVID-19 pandemic: position statements of the Asian Pacific Society for Digestive Endoscopy (APSDE-COVID statements). *Gut.* 2020;69:991–6, <http://dx.doi.org/10.1136/gutjnl-2020-321185>.
5. Lui RN, Wong SH, Sánchez-Luna SA, et al. Overview of guidance for endoscopy during the coronavirus disease 2019 pandemic. *J Gastroenterol Hepatol.* 2020;35:749–59, <http://dx.doi.org/10.1111/jgh.15053>.
6. Repici A, Maselli R, Colombo M, et al. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc.* 2020;92:192–7, <http://dx.doi.org/10.1016/j.gie.2020.03.019>.
7. Ruan W, Fishman DS, Lerner G, et al. Changes in pediatric endoscopic practice during the coronavirus disease 2019 pandemic: results from an international survey. *Gastroenterology.* 2020;159:1547–50, <http://dx.doi.org/10.1053/j.gastro.2020.05.068>.
8. Comunicado Técnico Diario: Información internacional y nacional sobre nuevo coronavirus con corte al 26 de enero de 2021. SSA, Gobierno de México. [Accessed el 25 de enero 2021]. Available from: <https://coronavirus.gob.mx/2021/01/26/conferencia-26-de-enero/>.
9. Say DS, de Lorimier A, Lammers CR, et al. Addendum to: risk stratification and personal protective equipment use in pediatric endoscopy during the COVID-19 outbreak: a single-center protocol. *JPGN.* 2020;70(6):755–6, <http://dx.doi.org/10.1097/MPG.0000000000002762>.
10. Gastroenterological Society of Australia (GESA) Devereaux B, Kaffes A, Strasser S, et al. Updated advice on preventative measures during gastrointestinal (GI) endoscopic procedures during the COVID-19 pandemic. *GESA.* 2020;1:1–9.
11. Walsh CM, Fishman DS, Lerner DG, et al. Pediatric endoscopy in the era of coronavirus disease 2019: A North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper. *JPGN.* 2020;70:741–50, <http://dx.doi.org/10.1097/MPG.0000000000002750>.